

CLEANING LABORATORY EVALUATION SUMMARY

SCL #: 2007
 DateRun: 09/17/2007
 Experimenters: Jason Marshall
 ClientType: Optical Manufacturer
 ProjectNumber: Project #1
 Substrates: Glass/Quartz
 PartType: Part
 Contaminants:
 Cleaning Methods: Immersion/Soak
 Analytical Methods: Gravimetric

Purpose: To evaluate supplied parts for resistance to alkaline solution.

Experimental Procedure: Glass samples were delivered polished to the specified polishing conditions. Each glass sample was weighed using an analytical balance. One sample of each glass type was hung by platinum wire into 0.01 mol/liter sodium hydroxide at 50 deg C for the length of times specified (15 minutes, 1 hour, 4 hours or 16 hours). After this treatment, the samples were weighed again and the loss of the mass of the sample was calculated. Calculation of the time t_{0.1} in hours, necessary to etch a surface layer a depth of 0.1 um was done using the following formula:

$$t_{0.1} = (t_e d S) / [(m_1 - m_2) * 100]$$

where:

t_{0.1} = the time (min) necessary to etch a surface layer to a depth of 0.1 μm

t_e = the time (min) for attach in the experiment

d = the specific gravity of the sample

S = the surface area (cm²) of the sample

m₁ = the mass (mg) of the sample before the test

m₂ = the mass (mg) of the sample after the test

Phosphate Resistance Class PR 1 2 3 4

Time t_{0.1} needed to etch to a depth of 0.1um/min >240 240~60 60~15 <15

In addition, changes in the surface of the sample following the treatment are qualitatively evaluated with the naked eye. Additional classification numbers are given according to the second table.

Additional Number Changes in the Surface

- .0 No visible changes
- .1 Clear, but irregular surface (wavy, pockmarked)
- .2 Interference colors (slight selective leaching)
- .3 Tenacious thin whitish layer (stronger selective)
- .4 Loosely adhering thick layer (surface crust)

Results: The target weight loss of 0.0010 grams (1 mg) was reached by two of the eight glass pieces (Sample A and Sample 1) after soaking for 16 hours. Samples 2 and B both were close to the 0.0010-gram level but did not exceed the value. The tables list the amount of weight loss after each time interval.

0.25 hour

| sample id | m1 | m2 | m1-m2 |
|-----------|--------|--------|---------|
| A | 3.806 | 3.8058 | 0.0002 |
| B | 3.8679 | 3.8679 | 0.0000 |
| C | 3.8306 | 3.8305 | 0.0001 |
| D | 4.1472 | 4.1472 | 0.0000 |
| 1 | 6.4285 | 6.4286 | -0.0001 |
| 2 | 1.2379 | 1.2379 | 0.0000 |
| 3 | 1.2177 | 1.2177 | 0.0000 |
| 4 | 1.2546 | 1.2543 | 0.0003 |

1 hour

| sample id | m1 | m2 | m1-m2 |
|-----------|--------|--------|---------|
| A | 3.806 | 3.8058 | 0.0002 |
| B | 3.8679 | 3.8675 | 0.0004 |
| C | 3.8306 | 3.8303 | 0.0003 |
| D | 4.1472 | 4.1467 | 0.0005 |
| 1 | 6.4285 | 6.4286 | -0.0001 |

CLEANING LABORATORY EVALUATION SUMMARY

| | | | |
|-----------|--------|--------|--------|
| 2 | 1.2379 | 1.2374 | 0.0005 |
| 3 | 1.2177 | 1.2177 | 0.0000 |
| 4 | 1.2546 | 1.2543 | 0.0003 |
| 4 hour | | | |
| sample id | m1 | m2 | m1-m2 |
| A | 3.806 | 3.8054 | 0.0006 |
| B | 3.8679 | 3.8671 | 0.0008 |
| C | 3.8306 | 3.8303 | 0.0003 |
| D | 4.1472 | 4.1464 | 0.0008 |
| 1 | 6.4285 | 6.4278 | 0.0007 |
| 2 | 1.2379 | 1.2372 | 0.0007 |
| 3 | 1.2177 | 1.2177 | 0.0000 |
| 4 | 1.2546 | 1.2543 | 0.0003 |

| Sample ID | density(d) | surface area(s) | d*S | time (te) | te*d*S | te*d*S/ [(m1-m2)*100] |
|-----------|------------|-----------------|---------|-----------|----------|-----------------------|
| A | 3.22 | 45.66 | 147.032 | 16 | 2352.506 | 23525.0624 |
| 1 | 2.23 | 53.9 | 120.197 | 16 | 1923.152 | 16026.2667 |

| Sample ID | Level |
|-----------|-------|
| A | AR1.0 |
| B | AR1.0 |
| C | AR1.0 |
| D | AR1.0 |
| 1 | AR1.0 |
| 2 | AR1.0 |
| 3 | AR1.0 |
| 4 | AR1.0 |

Summary:

Conclusion:

Six samples did not exceed the 1 mg weight loss. However, all eight samples would be classified as AR1.0 as the two samples that did loose over the 1 mg did so only after 16 hours of soaking. None of the samples had any visible abnormalities.